



Section: Pushdown Automata

Ch. 7 - Pushdown Automata

A DFA = $(Q, \Sigma, \delta, q_0, F)$



Modify DFA by adding a stack. New machine is called Pushdown Automata (PDA).



Definition: Nondeterministic PDA (NPDA) is defined by

$$M = (Q, \Sigma, \Gamma, \delta, q_0, z, F)$$

where

Q is finite set of states

Σ is tape (input) alphabet

Γ is stack alphabet

q_0 is initial state

z - start stack symbol, $z \in \Gamma$

$F \subseteq Q$ is set of final states.

$\delta: Q \times (\Sigma \cup \{\lambda\}) \times \Gamma \rightarrow$ finite subsets of $Q \times \Gamma^*$

Example of transitions

$$\delta(q_1, \mathbf{a}, \mathbf{b}) = \{(q_3, \mathbf{b}), (q_4, \mathbf{ab}), (q_6, \lambda)\}$$

The diagram for the above transitions is:

Instantaneous Description:

$$(q, w, u)$$

Description of a Move:

$$(q_1, aw, bx) \vdash (q_2, w, yx) \\ \text{iff}$$

Definition Let $M = (Q, \Sigma, \Gamma, \delta, q_0, z, F)$ be a NPDA. $L(M) = \{w \in \Sigma^* \mid (q_0, w, z) \vdash^* (p, \lambda, u), p \in F, u \in \Gamma^*\}$. The NPDA accepts all strings that start in q_0 and end in a final state.

Example: $\mathbf{L} = \{a^n b^n \mid n \geq 0\}$, $\Sigma = \{a, b\}$,
 $\Gamma = \{z, a\}$

Another Definition for Language Acceptance

NPDA M accepts $L(M)$ by empty stack:

$$L(M) = \{w \in \Sigma^* \mid (q_0, w, z) \vdash^* (p, \lambda, \lambda)\}$$

Example: $\mathbf{L} = \{a^n b^m c^{n+m} \mid n, m > 0\}$,
 $\Sigma = \{a, b, c\}$, $\Gamma = \{0, z\}$

Examples for you to try on your own:
(solutions are at the end of the
handout).

- $\mathbf{L} = \{a^n b^m \mid m > n, m, n > 0\}$, $\Sigma = \{a, b\}$,
 $\Gamma = \{z, a\}$
- $\mathbf{L} = \{a^n b^{n+m} c^m \mid n, m > 0\}$, $\Sigma = \{a, b, c\}$,
- $\mathbf{L} = \{a^n b^{2n} \mid n > 0\}$, $\Sigma = \{a, b\}$

Definition: A PDA

$M=(Q,\Sigma,\Gamma,\delta,q_0,z,F)$ is *deterministic* if
for every $q \in Q$, $a \in \Sigma \cup \{\lambda\}$, $b \in \Gamma$


1. $\delta(q, a, b)$ contains at most 1 element
2. if $\delta(q, \lambda, b) \neq \emptyset$ then $\delta(q, c, b)=\emptyset$ for all $c \in \Sigma$

Definition: L is DCFL iff \exists DPDA M
s.t. $L=L(M)$.


Examples:

1. Previous pda for $\{a^n b^n | n \geq 0\}$ is deterministic?
2. Previous pda for $\{a^n b^m c^{n+m} | n, m > 0\}$ is deterministic?
3. Previous pda for $\{ww^R | w \in \Sigma^+\}, \Sigma = \{a, b\}$ is deterministic?

Example: $L = \{a^n b^m \mid m > n, m, n > 0\}$,
 $\Sigma = \{a, b\}$, $\Gamma = \{z, a\}$



Example: $L = \{a^n b^{n+m} c^m \mid n, m > 0\}$,
 $\Sigma = \{a, b, c\}$,



Example: $L = \{a^n b^{2n} \mid n > 0\}$, $\Sigma = \{a, b\}$

