

Which of the following languages are CFL?

- $L = \{a^n b^n c^j \mid 0 < n \leq j\}$
- $L = \{a^n b^j a^n b^j \mid n > 0, j > 0\}$
- $L = \{a^n b^j a^k b^p \mid n + j \leq k + p, n > 0, j > 0, k > 0, p > 0\}$
- $L = \{a^n b^j a^j b^n \mid n > 0, j > 0\}$

**Pumping Lemma for Regular Language's:** Let  $L$  be a regular language, Then there is a constant  $m$  such that  $w \in L$ ,  $|w| \geq m$ ,  $w = xyz$  such that

- $|xy| \leq m$
- $|y| \geq 1$
- for all  $i \geq 0$ ,  $xy^i z \in L$

**Pumping Lemma for CFL's** Let  $L$  be any infinite CFL. Then there is a constant  $m$  depending only on  $L$ , such that for every string  $w$  in  $L$ , with  $|w| \geq m$ , we may partition  $w = uvxyz$  such that:

$$\begin{aligned} &|vxy| \leq m, \text{ (limit on size of substring)} \\ &|vy| \geq 1, \text{ (} v \text{ and } y \text{ not both empty)} \\ &\text{For all } i \geq 0, uv^i xy^i z \in L \end{aligned}$$

- **Proof:** (sketch) There is a CFG  $G$  s.t.  $L=L(G)$ .

Consider the parse tree of a long string in  $L$ .

For any long string, some nonterminal  $N$  must appear twice in the path.

**Example:** Consider  $L = \{a^n b^n c^n : n \geq 1\}$ . Show  $L$  is not a CFL.

• **Proof:** (by contradiction)

Assume  $L$  is a CFL and apply the pumping lemma.

Let  $m$  be the constant in the pumping lemma and consider  $w = a^m b^m c^m$ . Note  $|w| \geq m$ .

Show there is no division of  $w$  into  $uvxyz$  such that  $|vy| \geq 1$ ,  $|vxy| \leq m$ , and  $uv^i xy^i z \in L$  for  $i = 0, 1, 2, \dots$

Case 1: Neither  $v$  nor  $y$  can contain 2 or more distinct symbols. If  $v$  contains  $a$ 's and  $b$ 's, then  $uv^2 xy^2 z \notin L$  since there will be  $b$ 's before  $a$ 's.

Thus,  $v$  and  $y$  can be only  $a$ 's,  $b$ 's, or  $c$ 's (not mixed).

Case 2:  $v = a^{t_1}$ , then  $y = a^{t_2}$  or  $b^{t_3}$  ( $|vxy| \leq m$ )

If  $y = a^{t_2}$ , then  $uv^2 xy^2 z = a^{m+t_1+t_2} b^m c^m \notin L$  since  $t_1 + t_2 > 0$ ,  $n(a) > n(b)$ 's (number of  $a$ 's is greater than number of  $b$ 's)

If  $y = b^{t_3}$ , then  $uv^2 xy^2 z = a^{m+t_1} b^{m+t_3} c^m \notin L$  since  $t_1 + t_3 > 0$ , either  $n(a) > n(c)$ 's or  $n(b) > n(c)$ 's.

Case 3:  $v = b^{t_1}$ , then  $y = b^{t_2}$  or  $c^{t_3}$

If  $y = b^{t_2}$ , then  $uv^2 xy^2 z = a^m b^{m+t_1+t_2} c^m \notin L$  since  $t_1 + t_2 > 0$ ,  $n(b) > n(a)$ 's.

If  $y = c^{t_3}$ , then  $uv^2 xy^2 z = a^m b^{m+t_1} c^{m+t_3} \notin L$  since  $t_1 + t_3 > 0$ , either  $n(b) > n(a)$ 's or  $n(c) > n(a)$ 's.

Case 4:  $v = c^{t_1}$ , then  $y = c^{t_2}$

then,  $uv^2 xy^2 z = a^m b^m c^{m+t_1+t_2} \notin L$  since  $t_1 + t_2 > 0$ ,  $n(c) > n(a)$ 's.

Thus, there is no breakdown of  $w$  into  $uvxyz$  such that  $|vy| \geq 1$ ,  $|vxy| \leq m$  and for all  $i \geq 0$ ,  $uv^i xy^i z$  is in  $L$ . Contradiction, thus,  $L$  is not a CFL. Q.E.D.

**Example** Why would we want to recognize a language of the type  $\{a^n b^n c^n : n \geq 1\}$ ?

**Example:** Consider  $L = \{a^n b^n c^p : p > n > 0\}$ . Show  $L$  is not a CFL.

- **Proof:** Assume  $L$  is a CFL and apply the pumping lemma. Let  $m$  be the constant in the pumping lemma and consider  $w = \underline{\hspace{2cm}}$ . Note  $|w| \geq m$ .  
Show there is no division of  $w$  into  $uvxyz$  such that  $|vy| \geq 1$ ,  $|vxy| \leq m$ , and  $uw^i xy^i z \in L$  for  $i = 0, 1, 2, \dots$

Thus, there is no breakdown of  $w$  into  $uvxyz$  such that  $|vy| \geq 1$ ,  $|vxy| \leq m$  and for all  $i \geq 0$ ,  $uw^i xy^i z$  is in  $L$ . Contradiction, thus,  $L$  is not a CFL. Q.E.D.

**Example:** Consider  $L = \{a^j b^k : k = j^2\}$ . Show  $L$  is not a CFL.

- **Proof:** Assume  $L$  is a CFL and apply the pumping lemma. Let  $m$  be the constant in the pumping lemma and consider  $w = \underline{\hspace{2cm}}$

Show there is no division of  $w$  into  $uvxyz$  such that  $|vy| \geq 1$ ,  $|vxy| \leq m$ , and  $uv^i xy^i z \in L$  for  $i = 0, 1, 2, \dots$

Case 1: Neither  $v$  nor  $y$  can contain 2 or more distinct symbols. If  $v$  contains  $a$ 's and  $b$ 's, then  $uv^2 xy^2 z \notin L$  since there will be  $b$ 's before  $a$ 's.

Thus,  $v$  and  $y$  can be only  $a$ 's, and  $b$ 's (not mixed).

Thus, there is no breakdown of  $w$  into  $uvxyz$  such that  $|vy| \geq 1$ ,  $|vxy| \leq m$  and for all  $i \geq 0$ ,  $uv^i xy^i z$  is in  $L$ . Contradiction, thus,  $L$  is not a CFL. Q.E.D.

**Exercise:** Prove the following is not a CFL by applying the pumping lemma. (answer is at the end of this handout).

Consider  $L = \{a^{2n} b^{2p} c^n d^p : n, p \geq 0\}$ . Show  $L$  is not a CFL.

**Example:** Consider  $L = \{w\bar{w}w : w \in \Sigma^*\}$ ,  $\Sigma = \{a, b\}$ , where  $\bar{w}$  is the string  $w$  with each occurrence of  $a$  replaced by  $b$  and each occurrence of  $b$  replaced by  $a$ . For example,  $w = baaa$ ,  $\bar{w} = abbb$ ,  $w\bar{w} = baaaabbb$ . Show  $L$  is not a CFL.

- **Proof:** Assume  $L$  is a CFL and apply the pumping lemma. Let  $m$  be the constant in the pumping lemma and consider  $w = \underline{\hspace{2cm}}$   
Show there is no division of  $w$  into  $uvxyz$  such that  $|vy| \geq 1$ ,  $|vxy| \leq m$ , and  $uv^i xy^i z \in L$  for  $i = 0, 1, 2, \dots$

Thus, there is no breakdown of  $w$  into  $uvxyz$  such that  $|vy| \geq 1$ ,  $|vxy| \leq m$  and for all  $i \geq 0$ ,  $uv^i xy^i z$  is in  $L$ . Contradiction, thus,  $L$  is not a CFL. Q.E.D.

**Example:** Consider  $L = \{a^n b^p b^p a^n\}$ .  $L$  is a CFL. The pumping lemma should apply!

Let  $m \geq 4$  be the constant in the pumping lemma. Consider  $w = a^m b^m b^m a^m$ .

We can break  $w$  into  $uvxyz$ , with:

If you apply the pumping lemma to a CFL, then you should find a partition of  $w$  that works!

## Chap 8.2 Closure Properties of CFL's

**Theorem** CFL's are closed under union, concatenation, and star-closure.

- **Proof:**

Given 2 CFG  $G_1 = (V_1, T_1, S_1, P_1)$  and  $G_2 = (V_2, T_2, S_2, P_2)$

- Union:

Construct  $G_3$  s.t.  $L(G_3) = L(G_1) \cup L(G_2)$ .

$G_3 = (V_3, T_3, S_3, P_3)$

- Concatenation:

Construct  $G_3$  s.t.  $L(G_3) = L(G_1) \circ L(G_2)$ .

$G_3 = (V_3, T_3, S_3, P_3)$

- Star-Closure  
Construct  $G_3$  s.t.  $L(G_3) = L(G_1)^*$   
 $G_3 = (V_3, T_3, S_3, P_3)$

QED.

**Theorem** CFL's are NOT closed under intersection and complementation.

- **Proof:**

- Intersection:

- Complementation:

**Theorem:** CFL's are closed under *regular* intersection. If  $L_1$  is CFL and  $L_2$  is regular, then  $L_1 \cap L_2$  is CFL.

- **Proof:** (sketch) This proof is similar to the construction proof in which we showed regular languages are closed under intersection. We take a NPDA for  $L_1$  and a DFA for  $L_2$  and construct a NPDA for  $L_1 \cap L_2$ .

$M_1 = (Q_1, \Sigma, \Gamma, \delta_1, q_0, z, F_1)$  is an NPDA such that  $L(M_1) = L_1$ .

$M_2 = (Q_2, \Sigma, \delta_2, q'_0, F_2)$  is a DFA such that  $L(M_2) = L_2$ .

Example of replacing arcs (NOT a Proof!):



Note this is not a proof, but sketches how we will combine the DFA and NPDA. We must formally define  $\delta_3$ . If

then

Must show

if and only if

Must show:

$w \in L(M_3)$  iff  $w \in L(M_1)$  and  $w \in L(M_2)$ .

QED.

## Questions about CFL:

1. Decide if CFL is empty?

2. Decide if CFL is infinite?

**Example:** Consider  $L = \{a^{2n}b^{2m}c^nd^m : n, m \geq 0\}$ . Show  $L$  is not a CFL.

- **Proof:** Assume  $L$  is a CFL and apply the pumping lemma. Let  $m$  be the constant in the pumping lemma and consider  $w = a^{2m}b^{2m}c^md^m$ .

Show there is no division of  $w$  into  $uvxyz$  such that  $|vy| \geq 1$ ,  $|vxy| \leq m$ , and  $uv^ixy^iz \in L$  for  $i = 0, 1, 2, \dots$

**Case 1:** Neither  $v$  nor  $y$  can contain 2 or more distinct symbols. If  $v$  contains  $a$ 's and  $b$ 's, then  $uv^2xy^2z \notin L$  since there will be  $b$ 's before  $a$ 's.

Thus,  $v$  and  $y$  can be only  $a$ 's,  $b$ 's,  $c$ 's, or  $d$ 's (not mixed).

**Case 2:**  $v = a^{t_1}$ , then  $y = a^{t_2}$  or  $b^{t_3}$  ( $|vxy| \leq m$ )

If  $y = a^{t_2}$ , then  $uv^2xy^2z = a^{2m+t_1+t_2}b^{2m}c^md^m \notin L$  since  $t_1 + t_2 > 0$ , the number of  $a$ 's is not twice the number of  $c$ 's.

If  $y = b^{t_3}$ , then  $uv^2xy^2z = a^{2m+t_1}b^{2m+t_3}c^md^m \notin L$  since  $t_1 + t_3 > 0$ , either the number of  $a$ 's (denoted  $n(a)$ ) is not twice  $n(c)$  or  $n(b)$  is not twice  $n(d)$ .

**Case 3:**  $v = b^{t_1}$ , then  $y = b^{t_2}$  or  $c^{t_3}$

If  $y = b^{t_2}$ , then  $uv^2xy^2z = a^{2m}b^{2m+t_1+t_2}c^md^m \notin L$  since  $t_1 + t_2 > 0$ ,  $n(b) > 2*n(d)$ .

If  $y = c^{t_3}$ , then  $uv^2xy^2z = a^{2m}b^{2m+t_1}c^{m+t_3}d^m \notin L$  since  $t_1 + t_3 > 0$ , either  $n(b) > 2*n(d)$  or  $2*n(c) > n(a)$ .

**Case 4:**  $v = c^{t_1}$ , then  $y = c^{t_2}$  or  $d^{t_3}$

If  $y = c^{t_2}$ , then  $uv^2xy^2z = a^{2m}b^{2m}c^{m+t_1+t_2}d^m \notin L$  since  $t_1 + t_2 > 0$ ,  $2*n(c) > n(a)$ .

If  $y = d^{t_3}$ , then  $uv^2xy^2z = a^{2m}b^{2m}c^{m+t_1}d^{m+t_3} \notin L$  since  $t_1 + t_3 > 0$ , either  $2*n(c) > n(a)$  or  $2*n(d) > n(b)$ .

**Case 5:**  $v = d^{t_1}$ , then  $y = d^{t_2}$

then  $uv^2xy^2z = a^{2m}b^{2m}c^md^{m+t_1+t_2} \notin L$  since  $t_1 + t_2 > 0$ ,  $2*n(d) > n(c)$ .

Thus, there is no breakdown of  $w$  into  $uvxyz$  such that  $|vy| \geq 1$ ,  $|vxy| \leq m$  and for all  $i \geq 0$ ,  $uv^ixy^iz$  is in  $L$ . Contradiction, thus,  $L$  is not a CFL. Q.E.D.